

ALGEBRAIC CURVATURE TENSORS WHOSE SKEW-SYMMETRIC CURVATURE OPERATOR HAS CONSTANT RANK 2

PETER GILKEY AND TAN ZHANG

ABSTRACT. Let R be an algebraic curvature tensor for a non-degenerate inner product of signature (p, q) where $q \geq 5$. If π is a spacelike 2 plane, let $R(\pi)$ be the associated skew-symmetric curvature operator. We classify the algebraic curvature tensors so $R(\cdot)$ has constant rank 2 and show these are geometrically realizable by hypersurfaces in flat spaces. We also classify the Ivanov-Petrova algebraic curvature tensors of rank 2; these are the algebraic curvature tensors of constant rank 2 such that the complex Jordan normal form of $R(\cdot)$ is constant.

§1 INTRODUCTION

Let ${}^g\nabla$ be the Levi-Civita connection of a pseudo-Riemannian manifold (M, g) of signature (p, q) . Let ${}^gR(x, y) := {}^g\nabla_x {}^g\nabla_y - {}^g\nabla_y {}^g\nabla_x - {}^g\nabla_{[x, y]}$ be the Riemann curvature operator and let ${}^gR(x, y, z, w)$ be the associated curvature tensor. We have:

$$\begin{aligned} (1.1.a) \quad & g({}^gR(x, y)z, w) = {}^gR(x, y, z, w), \\ (1.1.b) \quad & {}^gR(x, y, z, w) = -{}^gR(y, x, z, w) = -{}^gR(x, y, w, z), \\ (1.1.c) \quad & {}^gR(x, y, z, w) = {}^gR(z, w, x, y), \text{ and} \\ (1.1.d) \quad & {}^gR(x, y, z, w) + {}^gR(y, z, x, w) + {}^gR(z, x, y, w) = 0. \end{aligned}$$

It is convenient to work in a purely algebraic context. Let $\langle \cdot, \cdot \rangle$ be a non-degenerate symmetric inner product of signature (p, q) on a finite dimensional real vector space V . We say that a 4 tensor $R \in \otimes^4(V^*)$ is an *algebraic curvature tensor* if R satisfies the identities of equations (1.1.b), (1.1.c), and (1.1.d); the *associated algebraic curvature operator* $R(\cdot, \cdot)$ is then defined by equation (1.1.a). Given an inner product $\langle \cdot, \cdot \rangle_P$ and an algebraic curvature tensor R_P on the tangent space $T_P M$ at a point $P \in M$, there is the germ of a metric g on M so $g|_{T_P M} = \langle \cdot, \cdot \rangle_P$ and so ${}^gR|_{T_P M} = R_P$. Thus algebraic curvature tensors are important in differential geometry.

A central problem in differential geometry is to understand the relationship between algebraic properties of the curvature tensor gR and the underlying geometry of the manifold. The full curvature tensor is in general quite difficult to work with so one often uses the curvature tensor to define a natural endomorphism of the tangent bundle. One wants

1991 *Mathematics Subject Classification*. Primary 53B20.

Key words and phrases. Algebraic curvature tensor, skew-symmetric curvature operator, Lorentzian and higher signature, Ivanov-Petrova tensors.

to know the geometric consequences that follow if such an operator is assumed to have constant eigenvalues or to have constant rank. The Jacobi operator, the Stanilov operator, the Szabó operator, and the skew-symmetric curvature operator are examples of such natural operators. We refer to [6, 9] for further details; in the interests of brevity we shall content ourselves here with a brief discussion of the Jacobi and skew-symmetric curvature operators as motivation.

We say that a vector v is *spacelike* if $\langle v, v \rangle > 0$. We say that a subspace π of V is *spacelike* if the restriction of $\langle \cdot, \cdot \rangle$ to π is positive definite. Let

$$V^+ := \{v \in V : \langle v, v \rangle > 0\}, \quad \mathbb{S}^+(V) := \{v : \langle v, v \rangle = 1\}, \quad \text{and} \quad \text{Gr}_{(0,2)}^+(V, \langle \cdot, \cdot \rangle)$$

be the set of spacelike vectors, the set of unit spacelike vectors, and the set of oriented spacelike 2 planes in V . Let R be an algebraic curvature tensor. We define the *Jacobi operator* $J_R(x) : y \rightarrow R(y, x)x$. We say that R is *Osserman* if the complex eigenvalues of $J_R(\cdot)$ are constant on $\mathbb{S}^+(V)$. If (M, g) is a local rank 1 symmetric space or is flat, then the local isometries act transitively on the bundle $\mathbb{S}^+(TM)$ and hence the eigenvalues of J_R are constant on $\mathbb{S}^+(TM)$. In the Riemannian setting ($p = 0$), Osserman [14] wondered if the converse held; this question has become known as the Osserman conjecture. Chi [3] established the Osserman conjecture for Riemannian manifolds of dimension m for $m \equiv 1 \pmod{2}$, for $m \equiv 2 \pmod{4}$, and for $m = 4$. The corresponding conjecture for algebraic curvature tensors fails if $m \equiv 0 \pmod{4}$ [7]. In the Lorentzian setting ($p = 1$), the Osserman conjecture has been established [1, 4]. The Osserman conjecture is false in the higher signature setting ($p > 1$) [2, 5].

Let R be an algebraic curvature tensor. If $\{v_1, v_2\}$ is an oriented basis for a spacelike 2 plane π , the *skew-symmetric curvature operator* is defined independently of the basis by:

$$(1.1.e) \quad R(\pi) := \{\langle v_1, v_1 \rangle \langle v_2, v_2 \rangle - \langle v_1, v_2 \rangle^2\}^{-1/2} R(v_1, v_2).$$

We say R has *rank* r if $\text{Rank}(R(\pi)) = r$ for any oriented spacelike 2 plane π . The following theorem was proved using topological methods [10, 15]. It shows that $r = 2$ in many cases.

1.1 Theorem. *Let R be an algebraic curvature tensor of rank r .*

- (1) *Let $p \leq 1$. Let $q = 5$, $q = 6$, or $q \geq 9$. Then $r = 2$.*
- (2) *Let $p = 2$. Let $q \geq 10$. Assume neither q nor $q + 2$ are powers of 2. Then $r = 2$.*

If $(p, q) = (0, 4)$, if $(p, q) = (2, 2)$, or if $(p, q) = (4, 0)$, then the algebraic curvature tensors of rank 4 form an open non-empty subset of the space of all algebraic curvature tensors so the 4 dimensional setting is exceptional. It is not known if there exist algebraic curvature tensors of rank greater than 2 for other values of (p, q) . Kowalski et al [13] has shown in the Riemannian setting that an algebraic curvature tensor of rank 4 in dimension 4 must have both positive and negative sectional curvatures.

In light of Theorem 1.1, we shall focus our attention on the algebraic curvature tensors of rank 2. If ϕ is a linear map from V to V , we define a 4 tensor R_ϕ and associated operator by twisting the tensor of constant sectional curvature $+1$:

$$R_\phi(x, y)z := \langle \phi y, z \rangle \phi x - \langle \phi x, z \rangle \phi y, \quad \text{and} \\ R_\phi(x, y, z, w) := \langle \phi y, z \rangle \langle \phi x, w \rangle - \langle \phi x, z \rangle \langle \phi y, w \rangle.$$

The adjoint is defined by the identity $\langle \phi x, y \rangle = \langle x, \phi^* y \rangle$. We say that ϕ is *admissible* if $\phi^* = \phi$ and if the kernel of ϕ contains no spacelike vectors. We can classify rank 2 algebraic curvature tensors:

1.2 Theorem. *Let $q \geq 5$.*

- (1) *If ϕ is admissible, then $\pm R_\phi$ is a rank 2 algebraic curvature tensor.*
- (2) *If R is a rank 2 algebraic curvature tensor, then $R = \pm R_\phi$ for some admissible ϕ .*

We say that an algebraic curvature tensor R of rank r is *Ivanov-Petrova* if the complex Jordan normal form of $R(\pi)$ is the same for every oriented spacelike 2 plane π ; these tensors were first classified for $m = 4$ and $p = 0$ by Ivanov and Petrova [12]. The following result generalizes results of [10, 15] from the Riemannian and Lorentzian settings to arbitrary signatures.

1.3 Theorem. *Let $q \geq 5$. A rank 2 algebraic curvature tensor R is Ivanov-Petrova if and only if $R = \pm R_\phi$ for some admissible ϕ where one of the following conditions holds:*

- (1) *There exists $C \neq 0$ so $\langle \phi v_1, \phi v_2 \rangle = C \langle v_1, v_2 \rangle$ for all $v_i \in V$.*
- (2) *The range of ϕ is totally isotropic, i.e. $\langle \phi v_1, \phi v_2 \rangle = 0$ for all $v_i \in V$.*

To prove Theorem 1.2, it is convenient to decouple the domain and range and to establish a slightly more general classification result. Let $\langle \cdot, \cdot \rangle_A$ and $\langle \cdot, \cdot \rangle_B$ be non-degenerate symmetric inner products of signatures (p_A, q_A) and (p_B, q_B) on A and B . Let

$$\mathfrak{so}(B, \langle \cdot, \cdot \rangle_B) := \{\phi : \phi + \phi^* = 0\}$$

be the set of all skew-symmetric linear maps; this is the Lie algebra of the special orthogonal group defined by $\langle \cdot, \cdot \rangle_B$. Let T be an alternating bilinear map from $A \otimes A$ to $\mathfrak{so}(B, \langle \cdot, \cdot \rangle_B)$. The associated 4 tensor $T(a_1, a_2, b_1, b_2)$ defined by equation (1.1.a) has the symmetries of equation (1.1.b); conversely given a 4 tensor with the symmetries of equation (1.1.b), we can use equation (1.1.a) to define an alternating map from $A \otimes A$ to $\mathfrak{so}(B, \langle \cdot, \cdot \rangle_B)$. We use equation (1.1.e) to extend T to a map from $\text{Gr}_{(0,2)}^+(A, \langle \cdot, \cdot \rangle_A)$ to $\mathfrak{so}(B, \langle \cdot, \cdot \rangle_B)$ and suppose this extension has constant rank 2. Such a map will be said to be *admissible*.

Let ϕ be a linear map from A to B , let $\xi \in B$, and let χ be an alternating bilinear map from $A \otimes A$ to B . Define:

$$\begin{aligned} (1.3.a) \quad & T_{\chi, \xi}(a_1, a_2)b := \langle \chi(a_1, a_2), b \rangle_B \xi - \langle \xi, b \rangle_B \chi(a_1, a_2), \\ & T_{\chi, \xi}(a_1, a_2, b_1, b_2) := \langle \chi(a_1, a_2), b_1 \rangle_B \langle \xi, b_2 \rangle_B - \langle \xi, b_1 \rangle_B \langle \chi(a_1, a_2), b_2 \rangle_B, \\ & T_\phi(a_1, a_2)b := \langle \phi(a_2), b \rangle_B \phi(a_1) - \langle \phi(a_1), b \rangle_B \phi(a_2), \text{ and} \\ & T_\phi(a_1, a_2, b_1, b_2) = \langle \phi(a_2), b_1 \rangle_B \langle \phi(a_1), b_2 \rangle_B - \langle \phi(a_1), b_1 \rangle_B \langle \phi(a_2), b_2 \rangle_B. \end{aligned}$$

Since the symmetries of equation (1.1.b) are satisfied, $T_{\chi, \xi}$ and T_ϕ are alternating bilinear maps from $A \otimes A$ to $\mathfrak{so}(B, \langle \cdot, \cdot \rangle_B)$.

1.4 Theorem. *Let $q_A \geq 5$.*

- (1) *We have the following criteria for admissibility:*
 - 1-a) *T_ϕ is admissible if and only if $\ker(\phi) \cap A^+ = \emptyset$.*
 - 1-b) *$T_{\xi, \chi}$ is admissible if and only if $\{\chi(a_1, a_2), \xi\}$ are linearly independent vectors whenever $\{a_1, a_2\}$ spans a spacelike 2 plane.*
- (2) *Let T be admissible. Then either $T = T_{\chi, \xi}$ or $T = \pm T_\phi$ for suitable (χ, ξ) or ϕ .*

Let R be an algebraic curvature tensor. We say that a pseudo-Riemannian manifold (M, g) is a *geometric realization* of R at a point $P \in M$ if there is an isomorphism ψ from $T_P M$ to V so that $\psi^*\{\langle \cdot, \cdot \rangle_V\} = g(P)$ and so that $\psi^*\{R\} = {}^g R(P)$. It is a classical result that every algebraic curvature tensor has a geometric realization. Let R be a rank 2 algebraic curvature tensor. By Theorem 1.2, there exists an admissible ϕ so $R = \varepsilon R_\phi$ where $\varepsilon = \pm 1$. Let $e_{m+1} \cdot \mathbb{R}$ be a 1 dimensional vector space. Define:

$$(1.4.a) \quad \begin{aligned} W &:= V \oplus e_{m+1} \cdot \mathbb{R}, \quad F_\phi(v) := v \oplus \frac{1}{2} \langle \phi(v), v \rangle e_{m+1}, \text{ and} \\ \langle v_1 \oplus r_1 e_{m+1}, v_2 \oplus r_2 e_{m+1} \rangle_W &= \langle v_1, v_2 \rangle_V + \varepsilon r_1 r_2. \end{aligned}$$

Then F_ϕ defines the germ of an embedding of V into W as a hypersurface which is defined in some small neighborhood of the origin. Let g_ϕ be the associated first fundamental form. We take the canonical identification of $T_0 V = V$. We say that (M, g) has rank r if ${}^g R$ has rank r at each point of the manifold.

1.5 Theorem. *Let $R = \pm R_\phi$ be a rank 2 algebraic curvature tensor on $(V, \langle \cdot, \cdot \rangle_V)$. Then (V, g_ϕ) is the germ of a rank 2 pseudo-Riemannian manifold which realizes R at $P = 0$.*

We say that a pseudo-Riemannian manifold (M, g) is *Ivanov-Petrova* if (M, g) has constant rank r and if ${}^g R$ is Ivanov-Petrova at each point of M ; the complex Jordan normal form can vary with the point in question but the rank is assumed constant. The Riemannian Ivanov-Petrova manifolds have been classified if $m \geq 4$ and $m \neq 7$ [8, 10, 12]; the case $m = 7$ is exceptional and the classification is not complete although some partial results are known [11]. We refer to [15] for some partial results in the pseudo-Riemannian setting. This classification shows that there exist Ivanov-Petrova algebraic curvature tensors which are not geometrically realized by Ivanov-Petrova pseudo-Riemannian manifolds.

Here is a brief outline to the paper. In §2, we establish some technical results. In Lemma 2.1, we discuss linearizations of a projective map. In Lemmas 2.2, 2.3 and 2.4, we establish some of the elementary properties of $\mathfrak{so}(B, \langle \cdot, \cdot \rangle_B)$. In Lemma 2.6, we give necessary and sufficient conditions for T_ϕ to be an algebraic curvature tensor if $\ker(\phi)$ contains no spacelike vector. We conclude §2 by proving Theorems 1.2 (1) and 1.4 (1).

Let $\mathbb{P}(A)$ and $\mathbb{P}(B)$ be the projective spaces defined by A and B . Let $[a] = a \cdot \mathbb{R} \in \mathbb{P}(A)$ and $[b] := b \cdot \mathbb{R} \in \mathbb{P}(B)$ be the lines spanned by $0 \neq a \in A$ and $0 \neq b \in B$. Let T be admissible. We define

$$(1.5.a) \quad \Phi([a]) := \bigcap_{\pi \in \text{Gr}_{(0,2)}^+(A): [a] \subset \pi} \text{Range}(T(\pi)).$$

In §3, we assume $p_A = p_B = 0$. We show in Lemma 3.1 that $\dim \Phi = 1$. If Φ is constant, we choose $0 \neq \xi \in \Phi$, we set $\chi(a_1, a_2) := T(a_1, a_2)\xi$, and we show $T = C T_{\chi, \xi}$. If Φ is non-constant, we show there exists a map ϕ which linearizes Φ and we show $T = C T_\phi$ for some constant C . This completes the proof of Theorem 1.4 (2) in the Riemannian setting.

In §4, we assume $p_B = 0$ and let p_A be arbitrary. Let S be a maximal spacelike subspace of A and let $T_S = T|_{S \otimes S}$. We use the results of §3 to express $T_S = T_{\chi(S), \xi(S)}$ or $T_S = T_{\phi(S)}$. The main technical difficulty is then to extend these tensors to all of A and thereby prove Theorem 1.4 (2) in this setting. In §5, we use the map ψ defined in Lemma 2.2 (1) to show that the signature of the metric on B plays no role and to complete the proof of Theorem 1.4 (2). We then use Theorem 1.4 (2) to complete the proof of Theorem 1.2 (2). In §6, we use Theorem 1.2 to prove Theorem 1.3.

In §7, we review some facts concerning the geometry of hypersurfaces and use Theorem 1.2 to prove Theorem 1.5. In §8, we construct some examples. In Example 8.1, we show that the assumption $q_A \geq 5$ in Theorem 1.4 is essential by constructing an admissible map T from $\mathbb{R}^{(p,4)} \otimes \mathbb{R}^{(p,4)}$ to $\mathfrak{so}(4)$ so that $T \neq \pm T_\phi$ and $T \neq T_{\chi,\xi}$ for any ϕ or (χ, ξ) . In Example 8.2, we give an algebraic curvature tensor which has constant eigenvalues and constant rank 2 on the set of oriented spacelike 2 planes but which is not Ivanov-Petrova; this shows that it is necessary to consider the complex Jordan normal form and not simply the eigenspace structure and rank. A subspace π is said to be *timelike* if the restriction of $\langle \cdot, \cdot \rangle$ to π is negative definite. By replacing $\langle \cdot, \cdot \rangle$ by $-\langle \cdot, \cdot \rangle$ we can interchange the roles of timelike and spacelike subspaces. We use equation (1.1.e) to define $R(\pi)$ for oriented timelike 2 planes. In Example 8.3, we exhibit a rank 2 Ivanov-Petrova algebraic curvature tensor which does not have constant rank on the set of timelike 2 planes. Thus the behavior for spacelike and timelike planes can be different.

It is a pleasant task to acknowledge helpful conversations with Dr. Inis Kath dealing with stylistic matters in the paper.

§2 SOME TECHNICAL PRELIMINARIES

Let $\mathbb{P}(A^+) \subseteq \mathbb{P}(A)$ be the subset of spacelike lines in A . Let Φ be a map from $\mathbb{P}(A^+)$ to $\mathbb{P}(B)$. We say that a linear map ϕ from A to B *linearizes* Φ if $\ker(\phi)$ contains no spacelike vectors and if $[\phi(a)] = \Phi[a]$ for all $[a] \in \mathbb{P}(A^+)$. Note that vectors $\{a_1, \dots, a_p\}$ are linearly independent if and only if $a_1 \wedge \dots \wedge a_p \neq 0$ in $\Lambda^p(A)$.

2.1 Lemma. *Let $\Phi : \mathbb{P}(A^+) \rightarrow \mathbb{P}(B)$.*

- (1) *If $q_A \geq 3$ and if ϕ_i are linearizations of Φ , then ϕ_1 is a multiple of ϕ_2 .*
- (2) *Let $p_A = 0$ and let $q_A \geq 5$. Let $\mathcal{F}$ be a family of codimension 1 subspaces of A . Suppose Φ is linearizable on every $F \in \mathcal{F}$ and that every 2 plane is contained in some element of $\mathcal{F}$. Then Φ is linearizable.*

Proof. The spacelike vectors form an open subset of A which generates A additively. Thus to prove $\phi_1 = c\phi_2$ it suffices to prove that there exists c so $\phi_1(a) = c\phi_2(a)$ for every spacelike a . Let a_1 and a_2 be spacelike. Since $q_A \geq 3$, we can choose a_0 spacelike so $a_0 \perp a_1$ and $a_0 \perp a_2$. Since $[\phi_1 a_i] = [\phi_2 a_i]$, there are non-zero real numbers c_i so that $\phi_1(a_i) = c_i \phi_2(a_i)$. To prove assertion (1) it suffices to show $c_1 = c_0 = c_2$. Since the roles of a_1 and a_2 are symmetric, we must only show $c_0 = c_1$. Since ϕ_2 is injective on the spacelike 2 plane $\text{Span}\{a_0, a_1\}$, $0 \neq \phi_2(a_0) \wedge \phi_2(a_1)$. Choose c_{01} so that $\phi_1(a_0 + a_1) = c_{01} \phi_2(a_0 + a_1)$. We show $c_0 = c_{01} = c_1$ and complete the proof of assertion (1) by computing:

$$\begin{aligned} c_{01} \phi_2(a_0) + c_{01} \phi_2(a_1) &= c_{01} \phi_2(a_0 + a_1) = \phi_1(a_0 + a_1) \\ &= \phi_1(a_0) + \phi_1(a_1) = c_0 \phi_2(a_0) + c_1 \phi_2(a_1). \end{aligned}$$

Let ϕ_F be the linearization of Φ on $F \in \mathcal{F}$. Fix a basepoint $\bar{F} \in \mathcal{F}$. Both ϕ_F and $\phi_{\bar{F}}$ are linearizations of Φ on $F \cap \bar{F} \neq \{0\}$. Since $\dim(F \cap \bar{F}) \geq q_A - 2 \geq 3$, we can use assertion (1) to see $\phi_F = c(F)\phi_{\bar{F}}$ on $F \cap \bar{F}$. We replace ϕ_F by $c(F)^{-1}\phi_F$ to assume without loss of generality that $\phi_F = \phi_{\bar{F}}$ on $F \cap \bar{F}$ for all $F \in \mathcal{F}$. Let $F_i \in \mathcal{F}$. Since $\phi_{F_1} = \phi_{\bar{F}} = \phi_{F_2}$ on $F_1 \cap F_2 \cap \bar{F} \neq \{0\}$, we use assertion (1) to see $\phi_{F_1} = \phi_{F_2}$ on the (possibly) larger

intersection $F_1 \cap F_2$. Thus $\phi(a) = \phi_F(a)$ is well defined on $\cup_{F \in \mathcal{F}} F = A$. Since every 2 plane is contained in some element of $\mathcal{F}$, ϕ is linear and linearizes Φ . $\square$

We can relate $\langle \cdot, \cdot \rangle_B$ to a positive definite inner product.

2.2 Lemma. *There exists a self-adjoint linear map ψ from B to B with $\psi^2 = id$ so that $(b_1, b_2)_+ := \langle \psi b_1, b_2 \rangle_B$ is a positive definite symmetric inner product on B , so that $\langle b, \tilde{b} \rangle_B = (\psi b, \tilde{b})_+ = (b, \psi \tilde{b})_+$, and so that $(b, \tilde{b})_+ = \langle \psi b, \tilde{b} \rangle_B = \langle b, \psi \tilde{b} \rangle_B$. The map $\mathcal{T} \rightarrow \psi \mathcal{T}$ is an isomorphism between $\mathfrak{so}(B, \langle \cdot, \cdot \rangle_B)$ and $\mathfrak{so}(B, (\cdot, \cdot)_+)$.*

Proof. Choose a basis $\{\mathbf{b}_i\}$ for B so $\langle \mathbf{b}_i, \mathbf{b}_j \rangle_B = 0$ for $i \neq j$, so $\langle \mathbf{b}_i, \mathbf{b}_i \rangle_B = -1$ for $i \leq p_B$, and so $\langle \mathbf{b}_i, \mathbf{b}_i \rangle_B = 1$ for $i > p_B$. Let $\{\mathbf{b}^i\}$ be the corresponding dual basis for B^* . Define:

$$(b_1, b_2)_+ := \sum_i \mathbf{b}^i(b_1) \mathbf{b}^i(b_2) \text{ and } \psi(b_1) := -\sum_{i \leq p_B} \mathbf{b}^i(b_1) \mathbf{b}_i + \sum_{i > p_B} \mathbf{b}^i(b_1) \mathbf{b}_i.$$

The first identities now follow. We complete the proof by establishing the following chain of equivalent statements:

- (1) $\mathcal{T} \in \mathfrak{so}(B, \langle \cdot, \cdot \rangle_B)$.
- (2) $\langle \mathcal{T} b, \tilde{b} \rangle_B + \langle b, \mathcal{T} \tilde{b} \rangle_B = 0$ for all $b, \tilde{b} \in B$.
- (3) $(\psi \mathcal{T} b, \tilde{b})_+ + (b, \psi \mathcal{T} \tilde{b})_+ = 0$ for all $b, \tilde{b} \in B$.
- (4) $\psi \mathcal{T} \in \mathfrak{so}(B, (\cdot, \cdot)_+)$. $\square$

If $T \in \mathfrak{so}(B, \langle \cdot, \cdot \rangle_B)$, define $\omega(T) \in \Lambda^2(B^*)$ by $\omega(T)(b_1, b_2) := \langle T b_1, b_2 \rangle_B$. The correspondence $T \rightarrow \omega(T)$ identifies $\mathfrak{so}(B, \langle \cdot, \cdot \rangle_B)$ with $\Lambda^2(B^*)$.

2.3 Lemma. *Let $0 \neq \mathcal{T} \in \mathfrak{so}(B, \langle \cdot, \cdot \rangle_B)$. Then $\text{Rank}(\mathcal{T})$ is even. We have $\text{Rank}(\mathcal{T}) = 2$ if and only if $\omega(\mathcal{T}) \wedge \omega(\mathcal{T}) = 0$.*

Proof. Since $\text{Rank}(\mathcal{T}) = \text{Rank}(\psi \mathcal{T})$ and $\omega_{\langle \cdot, \cdot \rangle_B}(\mathcal{T}) = \omega_{(\cdot, \cdot)_+}(\psi \mathcal{T})$, we may suppose the metric is positive definite. Choose an orthonormal basis $\{\mathbf{b}_i\}$ for B and $\lambda_\mu > 0$ so

$$\mathcal{T} \mathbf{b}_j = 0 \text{ for } j > 2\ell \text{ and } \mathcal{T} \mathbf{b}_{2\mu} = \lambda_\mu \mathbf{b}_{2\mu-1}, \mathcal{T} \mathbf{b}_{2\mu-1} = -\lambda_\mu \mathbf{b}_{2\mu} \text{ for } \mu \leq \ell.$$

Thus $\text{Rank}(\mathcal{T}) = 2\ell$ is even. Let $\{\mathbf{b}_i^*\}$ be the dual basis for B^* . We use the following decomposition to complete the proof:

$$\omega(\mathcal{T}) = \sum_{1 \leq \mu \leq \ell} \lambda_\mu \mathbf{b}_{2\mu-1}^* \wedge \mathbf{b}_{2\mu}^*. \quad \square$$

2.4 Lemma. *Let $\mathcal{T}_i \in \mathfrak{so}(B, \langle \cdot, \cdot \rangle_B)$. Assume $\text{Rank}(\mathcal{T}_i) = 2$.*

- (1) *Then $\mathcal{T}_1$ is a multiple of $\mathcal{T}_2$ if and only if $\text{Range}(\mathcal{T}_1) = \text{Range}(\mathcal{T}_2)$.*
- (2) *Let $p_B = 0$. Let $0 \neq b \in \text{Range}(\mathcal{T}_1)$. Then $\{b, \mathcal{T}_1 b\}$ is an orthogonal basis for $\text{Range}(\mathcal{T}_1)$.*

Proof. We use Lemma 2.2 to assume the metric is positive definite in the proof of assertion (1). Since $\text{Rank}(\mathcal{T}_i) = 2$, $\mathcal{T}_i$ is a multiple of a 90 degree rotation in the 2 plane $\text{Range}(\mathcal{T}_i)$ and vanishes on $\text{Range}(\mathcal{T}_i)^\perp$. Assertions (1) and (2) now follow. $\square$

The following technical observation will prove useful in later sections.

2.5 Lemma. *Let T be a bilinear map from $A \otimes A$ to a vector space C . Assume that $T(a_1, a_2) = 0$ whenever a_1 and a_2 span a spacelike 2 plane. Then $T = 0$.*

Proof. The set of such tensors $a_1 \otimes a_2$ generates $A \otimes A$ additively. Since T is bilinear, we may conclude $T = 0$. $\square$

We continue our preparatory steps with a final technical Lemma.

2.6 Lemma. *Let ϕ be a linear map from A to B so $\ker(\phi)$ contains no spacelike vector. Let $q_A \geq 3$.*

- (1) *If $T_{\tilde{\phi}} = \pm T_{\phi}$, then $\tilde{\phi} = \pm \phi$.*
- (2) *Let $A = B = V$. Then T_{ϕ} is an algebraic curvature tensor if and only if $\phi = \phi^*$.*

Proof. Use equation (1.5.a) to define Φ . Let a_1 be spacelike. Since $q_A \geq 3$, we can choose a_2 and a_3 so $\text{Span}\{a_1, a_2, a_3\}$ is spacelike. Then $0 \neq \phi(a_1) \wedge \phi(a_2) \wedge \phi(a_3)$ so

$$\begin{aligned} \phi(a_1) \cdot \mathbb{R} &\subseteq \Phi([a_1]) \subseteq \text{Range}(T_{\phi}(a_1, a_2)) \cap \text{Range}(T_{\phi}(a_1, a_3)) \\ &\subseteq \text{Span}\{\phi(a_1), \phi(a_3)\} \cap \text{Span}\{\phi(a_1), \phi(a_2)\} \subseteq \phi(a_1) \cdot \mathbb{R}. \end{aligned}$$

Thus $\Phi([a_1]) = \phi(a_1) \cdot \mathbb{R}$ and ϕ linearizes Φ . If $T_{\phi} = \pm T_{\tilde{\phi}}$, then $\tilde{\phi}$ also linearizes $\Phi = \tilde{\Phi}$ so by Lemma 2.1 (1), $\tilde{\phi} = c\phi$. As $T_{c\phi} = c^2 T_{\phi}$, $c^2 = \pm 1$ so $c = \pm 1$ and assertion (1) follows.

Let $A = B = V$. Since T_{ϕ} defines an alternating bilinear map from $V \otimes V$ to $\mathfrak{so}(V, \langle \cdot, \cdot \rangle)$, the curvature symmetry (1.1.b) is immediate. We compute:

$$\begin{aligned} T_{\phi}(a_3, a_4, a_1, a_2) &= \langle \phi(a_4), a_1 \rangle \langle \phi(a_3), a_2 \rangle - \langle \phi(a_3), a_1 \rangle \langle \phi(a_4), a_2 \rangle \\ &= \langle \phi^*(a_1), a_4 \rangle \langle \phi^*(a_2), a_3 \rangle - \langle \phi^*(a_1), a_3 \rangle \langle \phi^*(a_2), a_4 \rangle \\ &= T_{\phi^*}(a_1, a_2, a_3, a_4). \end{aligned}$$

Thus symmetry of equation (1.1.c) is satisfied if and only if $T_{\phi} = T_{\phi^*}$ or equivalently using assertion (1) if and only if $\phi = \pm \phi^*$. If $\phi^* = \phi$, we verify that the Bianchi identities of equation (1.1.d) are satisfied by computing:

$$\begin{aligned} &T_{\phi}(x, y)z + T_{\phi}(y, z)x + T_{\phi}(z, x)y \\ &= \langle \phi y, z \rangle \phi x - \langle \phi x, z \rangle \phi y + \langle \phi z, x \rangle \phi y - \langle \phi y, x \rangle \phi z + \langle \phi x, y \rangle \phi z - \langle \phi z, y \rangle \phi x \\ &= (\langle \phi y, z \rangle - \langle \phi z, y \rangle) \phi x + (\langle \phi z, x \rangle - \langle \phi x, z \rangle) \phi y + (\langle \phi x, y \rangle - \langle \phi y, x \rangle) \phi z = 0. \end{aligned}$$

Conversely if the Bianchi identities are satisfied, we suppose that $\phi = -\phi^*$ and argue for a contradiction. We compute:

$$\begin{aligned} (2.6.a) \quad 0 &= T_{\phi}(a_1, a_2)a_3 + T_{\phi}(a_2, a_3)a_1 + T_{\phi}(a_3, a_1)a_2 \\ &= \langle \phi a_2, a_3 \rangle \phi a_1 + \langle \phi a_3, a_1 \rangle \phi a_2 + \langle \phi a_1, a_3 \rangle \phi a_3 \\ &\quad - \langle \phi a_1, a_3 \rangle \phi a_2 - \langle \phi a_2, a_1 \rangle \phi a_3 - \langle \phi a_3, a_2 \rangle \phi a_1 \\ &= 2\langle \phi a_2, a_3 \rangle \phi a_1 + 2\langle \phi a_3, a_1 \rangle \phi a_2 + 2\langle \phi a_1, a_2 \rangle \phi a_3. \end{aligned}$$

Fix a_1 spacelike. Then $\phi a_1 \neq 0$. Since $q_A \geq 3$, we may choose a_2 so $a_2 \perp \phi a_1$ and so $\{a_1, a_2\}$ is a spacelike orthonormal set. If $a_3 \perp \phi a_1$, since $\phi a_1 \neq 0$, we use equation (2.6.a) to see that $a_3 \perp \phi a_2$. Thus ϕa_2 is a multiple of ϕa_1 . This is not possible as ϕ is injective on spacelike subspaces. Consequently we have $\phi = \phi^*$. $\square$

2.7 The proof of Theorem 1.4 (1). We have $\text{Rank}(T_\phi) \leq 2$ and $\text{Rank}(T_{\chi,\xi}) \leq 2$. By Lemma 2.3, the rank is even. Thus T_ϕ has constant rank 2 if and only if $\{\phi(a_1), \phi(a_2)\}$ are linearly independent vectors whenever $\{a_1, a_2\}$ spans a spacelike 2 plane. This is equivalent to the condition that $\ker(\phi)$ contains no spacelike vector. Similarly $T_{\chi,\xi}$ has constant rank 2 if and only if $0 \neq \chi(a_1, a_2) \wedge \xi$ whenever $\{a_1, a_2\}$ spans a spacelike 2 plane. $\square$

2.8 The Proof of Theorem 1.2 (1). Suppose that $\phi = \phi^*$ and that $\ker(\phi)$ contains no spacelike vectors. By Theorem 1.4 (1), R_ϕ has constant rank 2. By Lemma 2.6 (2), R_ϕ is an algebraic curvature tensor. $\square$

§3 PROPERTIES OF THE MAP Φ

We suppose $p_A = p_B = 0$ and work in the Riemannian setting in this section. The following result concerning Φ is central to our investigation.

3.1 Lemma. *Let $p_A = p_B = 0$ and let T be admissible. Let $0 \neq a_1 \wedge a_2 \wedge a_3$.*

- (1) *Let $q_A \geq 3$. We have $\dim\{\text{Range}(T(a_1, a_2)) \cap \text{Range}(T(a_1, a_3))\} = 1$.*
- (2) *Let $q_A \geq 5$. Let Φ be defined by equation (1.5.a). We have $\dim(\Phi([a_1])) = 1$.*
- (3) *Let $q_A \geq 5$. We have $\Phi([a_1]) = \text{Range}(T(a_1, a_2)) \cap \text{Range}(T(a_1, a_3))$.*
- (4) *Let $q_A \geq 5$. The map $\Phi : \mathbb{P}(A) \rightarrow \mathbb{P}(B)$ is either injective or constant.*

Proof. We adapt arguments used to study IP algebraic curvature tensors in [10]. Let $\pi_{ij} := \text{Span}\{a_i, a_j\}$ and let $u := \dim\{\text{Range}(T(\pi_{12})) + \text{Range}(T(\pi_{13}))\}$. Then

$$\dim\{\text{Range}(T(\pi_{12})) \cap \text{Range}(T(\pi_{13}))\} = 4 - u.$$

We may estimate that $4 \geq u \geq \dim \text{Range}(T(\pi_{12})) = 2$. To prove assertion (1) we must show $u = 3$ i.e. that $u \neq 2$ and $u \neq 4$. If $u = 2$, then $\text{Range}(T(\pi_{12})) = \text{Range}(T(\pi_{13}))$ is a 2 dimensional space so $T(\pi_{12}) = cT(\pi_{13})$ by Lemma 2.4 (1). Thus $T(a_1, a_2 - ca_3) = 0$ which is false. This shows that $u \neq 2$. Let $\{\mathbf{b}_i\}$ be an orthonormal basis for B so

$$(3.1.a) \quad \begin{aligned} \text{Range}(T(\pi_{12})) &= \text{Span}\{\mathbf{b}_1, \mathbf{b}_2\}, \quad T(\pi_{13})\mathbf{b}_1 \in \text{Span}\{\mathbf{b}_1, \mathbf{b}_2, \mathbf{b}_3\}, \text{ and} \\ \text{Range}(T(\pi_{12})) + \text{Range}(T(\pi_{13})) &\subseteq \text{Span}\{\mathbf{b}_1, \mathbf{b}_2, \mathbf{b}_3, \mathbf{b}_4\}. \end{aligned}$$

Define $\mathcal{T}_{ij} \in \mathfrak{so}(B, \langle \cdot, \cdot \rangle_B)$ with $\omega(\mathcal{T}_{ij}) = \mathbf{b}_i^* \wedge \mathbf{b}_j^*$ by $\mathcal{T}_{ij} : \mathbf{b}_i \rightarrow \mathbf{b}_j, \mathbf{b}_j \rightarrow -\mathbf{b}_i$, and $\mathbf{b}_k \rightarrow 0$ for $k \neq i, j$. By rescaling T , we may assume $T(\pi_{12}) = \mathcal{T}_{12}$. We use equation (3.1.a) to expand:

$$\begin{aligned} T(\pi_{12}) &= \mathcal{T}_{12}, \quad \omega(T(\pi_{12})) = \mathbf{b}_1^* \wedge \mathbf{b}_2^*, \\ T(\pi_{13}) &= c_{12}\mathcal{T}_{12} + c_{13}\mathcal{T}_{13} + c_{23}\mathcal{T}_{23} + c_{24}\mathcal{T}_{24} + c_{34}\mathcal{T}_{34}, \text{ and} \\ \omega(T(\pi_{13})) &= c_{12}\mathbf{b}_1^* \wedge \mathbf{b}_2^* + c_{13}\mathbf{b}_1^* \wedge \mathbf{b}_3^* + c_{23}\mathbf{b}_2^* \wedge \mathbf{b}_3^* + c_{24}\mathbf{b}_2^* \wedge \mathbf{b}_4^* + c_{34}\mathbf{b}_3^* \wedge \mathbf{b}_4^*. \end{aligned}$$

Let $\mathcal{T}(\varepsilon) := T(a_1, a_2 + \varepsilon a_3)$. Since $\text{Rank}(\mathcal{T}(\varepsilon)) = 2$, we use Lemma 2.3 to see:

$$(3.1.b) \quad 0 = \omega(\mathcal{T}(\varepsilon)) \wedge \omega(\mathcal{T}(\varepsilon)) = \{2\varepsilon c_{34} + \varepsilon^2(c_{12}c_{34} - c_{13}c_{24})\}\mathbf{b}_1^* \wedge \mathbf{b}_2^* \wedge \mathbf{b}_3^* \wedge \mathbf{b}_4^*.$$

Since equation (3.1.b) holds for all $\varepsilon \in \mathbb{R}$, $c_{34} = 0$ and $c_{13}c_{24} = 0$. We complete the proof of assertion (1) by dealing with the two possible cases:

- (1) If $c_{24} = 0$, then $T(\pi_{13}) = c_{12}\mathcal{T}_{12} + c_{13}\mathcal{T}_{13} + c_{23}\mathcal{T}_{23}$ so $\text{Range}(T(\pi_{13})) \subseteq \text{Span}\{e_1, e_2, e_3\}$ and $u \leq 3$.
- (2) If $c_{13} = 0$, then $T(\pi_{13}) = c_{12}\mathcal{T}_{12} + c_{23}\mathcal{T}_{23} + c_{24}\mathcal{T}_{24}$ so $\text{Range}(T(\pi_{13})) \subseteq \text{Span}\{e_1, e_2, c_{23}e_3 + c_{24}e_4\}$ and $u \leq 3$.

Suppose assertion (2) fails; we argue for a contradiction. Choose a_4 with $0 \neq a_1 \wedge a_4$ so

$$(3.1.c) \quad \cap_{2 \leq i \leq 4} \text{Range}(T(a_1, a_i)) = \{0\}.$$

If $2 \leq i < j \leq 4$, then $\text{Range}(T(a_1, a_i)) \neq \text{Range}(T(a_1, a_j))$ so $0 \neq a_1 \wedge a_i \wedge a_j$. We use assertion (1) to define:

$$\begin{aligned} L_{ij} &:= \text{Range}(T(a_1, a_i)) \cap \text{Range}(T(a_1, a_j)) \in \mathbb{P}(B) \\ E &:= L_{23} + L_{24} + L_{34}. \end{aligned}$$

Let $\{i, j, k\}$ be a permutation of $\{2, 3, 4\}$. Then

$$\{0\} = \text{Range}(T(a_1, a_i)) \cap \text{Range}(T(a_1, a_j)) \cap \text{Range}(T(a_1, a_k)) = L_{ij} \cap L_{ik}.$$

Thus L_{ij} and L_{ik} are distinct lines which are contained in $\text{Range}(T(a_1, a_i))$. This shows:

$$\text{Range}(T(a_1, a_i)) = L_{ij} \oplus L_{ik} \text{ so } \text{Range}(T(a_1, a_i)) \subseteq E.$$

Let $[a_5] \perp [a_1]$. We wish to show $\text{Range}(T(a_1, a_5)) \subseteq E$. If $0 = a_1 \wedge a_i \wedge a_5$ for some i with $2 \leq i \leq 4$, then $T(a_1, a_5) = cT(a_1, a_i)$ so $\text{Range}(T(a_1, a_5)) \subseteq E$. Thus we suppose that $0 \neq a_1 \wedge a_i \wedge a_5$ for all $2 \leq i \leq 4$ and use assertion (1) to define the lines

$$L_{i5} := \text{Range}(T(a_1, a_i)) \cap \text{Range}(T(a_1, a_5)) \in \mathbb{P}(B) \text{ for } 2 \leq i \leq 4.$$

If $L_{25} = L_{35} = L_{45}$, then $L_{25} = L_{i5} \subseteq \text{Range}(T(a_1, a_i))$ for $2 \leq i \leq 4$ which contradicts equation (3.1.c). Thus at least two of the lines L_{i5} are distinct lines which are contained in $\text{Range}(T(a_1, a_5))$. Consequently the lines $\{L_{25}, L_{35}, L_{45}\}$ span $\text{Range}(T(a_1, a_5))$ and

$$\text{Range}(T(a_1, a_5)) \subseteq E.$$

This shows that the map $\psi : a_4 \rightarrow \text{Range}(T(a_1, a_4))$ is a well defined map from the projective space $\mathbb{P}(a_1^\perp)$ to the unoriented Grassmannian $\text{Gr}_2(E)$ of 2 planes in E . Suppose given distinct lines $[a_4]$ and $[a_5]$ in $\mathbb{P}(a_1^\perp)$. Since $0 \neq a_1 \wedge a_4 \wedge a_5$, we use assertion (1) to see $\psi([a_4]) \neq \psi([a_5])$. Thus ψ is injective. Since $\dim(E) \leq 3$, $\dim \text{Gr}_2(E) \leq 2$. It is immediate from the definition that ψ is continuous. We can therefore use invariance of domain to see $\dim(\mathbb{P}(a_1^\perp)) \leq 2$. Since $\dim(\mathbb{P}(a_1^\perp)) = q_A - 2 \geq 3$, this contradiction completes the proof of assertion (2). Assertions (3) now follow. In §8, we shall present an example that shows assertion (2) fails if $q_A = 4$.

Suppose that $\Phi : \mathbb{P}(A) \rightarrow \mathbb{P}(B)$ is not injective. Choose distinct lines $[a_i]$ so that $\Phi([a_1]) = \Phi([a_2])$. If $[a_3] \in \mathbb{P}(A) - \mathbb{P}(\text{Span}\{a_1, a_2\})$, then

$$\Phi([a_1]) = \Phi([a_2]) \subseteq \text{Range}(T(a_3, a_1)) \cap \text{Range}(T(a_3, a_2)) = \Phi([a_3]).$$

Thus Φ is constant on $\mathbb{P}(A) - \mathbb{P}(\text{Span}\{a_1, a_2\})$. We use assertion (2) to see Φ is continuous. It now follows that Φ is constant on all of $\mathbb{P}(A)$. $\square$

Theorem 1.4 (2) follows for positive definite metrics from the following Lemma:

3.2 Lemma. *Let $p_A = p_B = 0$ and let $q_A \geq 5$. Let T be admissible.*

- (1) *Let Φ be constant. Fix $\xi \in \mathbb{S}^+(\Phi)$. Set $\chi(a_1, a_2) := -T(a_1, a_2)\xi$. Then $T = T_{\chi, \xi}$.*
- (2) *Let Φ be non-constant. Then there exists a linearization ϕ of Φ so $T = \pm T_\phi$.*

Proof. Suppose $\Phi : \mathbb{P}(A) \rightarrow \mathbb{P}(B)$ is constant. Choose $\xi \in \mathbb{S}^+(\Phi)$. Then $\xi \in \Phi[a]$ for all a and we can define an alternating bilinear map χ from $A \otimes A$ to $\xi^\perp \subseteq B$ by setting $\chi(a_1, a_2) := -T(a_1, a_2)\xi \in B$. If $0 \neq a_1 \wedge a_2$, then we have

$$\text{Range}(T(a_1, a_2)) = \text{Span}\{\xi, \chi(a_1, a_2)\} = \text{Range}(T_{\chi, \xi}(a_1, a_2)).$$

Thus by Lemma 2.4 (1), $T(a_1, a_2) = cT_{\chi, \xi}(a_1, a_2)$. Since

$$T(a_1, a_2)\xi = -\chi(a_1, a_2) = T_{\chi, \xi}(a_1, a_2)\xi,$$

$c = 1$ so $T(a_1, a_2) = T_{\chi, \xi}(a_1, a_2)$ if $0 \neq a_1 \wedge a_2$. Lemma 2.5 now shows $T = T_{\chi, \xi}$ which completes the proof of assertion (1).

If Φ is not constant, then Φ is injective by Lemma 3.1 (4). The unit spheres $\mathbb{S}^+(A)$ and $\mathbb{S}^+(B)$ are the universal covers of $\mathbb{P}(A)$ and $\mathbb{P}(B)$. We lift Φ to define:

$$\Phi^+ : \mathbb{S}^+(A) \rightarrow \mathbb{S}^+(B) \text{ so } [\Phi^+(a)] = \Phi([a]).$$

Let $\{\mathfrak{a}_i\}$ be a basis for A and let $\{\mathfrak{a}^i\}$ be the corresponding dual basis for A^* . Let $i < j$. Since $\Phi([\mathfrak{a}_i])$ and $\Phi([\mathfrak{a}_j])$ are distinct lines which are contained in $T(\mathfrak{a}_i, \mathfrak{a}_j)$,

$$\text{Range}(T(\mathfrak{a}_i, \mathfrak{a}_j)) = \text{Span}\{\Phi^+(\mathfrak{a}_i), \Phi^+(\mathfrak{a}_j)\}.$$

Since T is bilinear, we have $T(a_1, a_2) = \sum_{i,j} \mathfrak{a}^i(a_1)\mathfrak{a}^j(a_2)T(\mathfrak{a}_i, \mathfrak{a}_j)$ so:

$$\text{Range}(T(a_1, a_2)) \subseteq \sum_{i,j} \text{Range}(T(\mathfrak{a}_i, \mathfrak{a}_j)) \subseteq \text{Span}\{\Phi^+(\mathfrak{a}_1), \dots, \Phi^+(\mathfrak{a}_{q_A})\}.$$

Thus replacing B by $\text{Span}\{\Phi^+(\mathfrak{a}_1), \dots, \Phi^+(\mathfrak{a}_{q_A})\}$ we may assume without loss of generality

$$\dim(B) \leq \dim(A).$$

Since $\mathbb{P}(A)$ is compact, $\text{Range}(\Phi)$ is compact and hence closed. We have that Φ is injective and that $\dim(\mathbb{P}(B)) \leq \dim(\mathbb{P}(A))$. Thus we use invariance of domain to see that $\dim(\mathbb{P}(B)) = \dim(\mathbb{P}(A))$ and that Φ is an open map. Thus $\text{Range}(\Phi)$ is an open subset of $\mathbb{P}(B)$. Since $\mathbb{P}(B)$ is connected and $\text{Range}(\Phi)$ is non-empty, Φ is surjective. Since $\mathbb{P}(A)$ is compact and since $\mathbb{P}(B)$ is Hausdorff, the fundamental theorem of point set topology shows that Φ is a homeomorphism from $\mathbb{P}(A)$ onto $\mathbb{P}(B)$. It now follows that the lift Φ^+ is a homeomorphism from $\mathbb{S}^+(A)$ onto $\mathbb{S}^+(B)$. Since Φ induces an isomorphism from the fundamental group of $\mathbb{P}(A)$ to the fundamental group of $\mathbb{P}(B)$, we have:

$$\Phi^+(-a) = -\Phi^+(a).$$

We now show that Φ is linearizable; such a linearization is unique up to constant multiple by Lemma 2.1 (1). Let $0 \neq a_1 \wedge a_2$. We first show $\Phi([a_1 + a_2]) \subseteq \Phi([a_1]) + \Phi([a_2])$. Since $q_B = q_A \geq 5$ and since Φ^+ is surjective, we choose unit vectors a_3 and a_4 so that

$$(3.2.a) \quad \Phi^+(a_3) \perp \Phi^+(a_4) \text{ and } \Phi^+(a_3) \perp \{\Phi^+(a_1), \Phi^+(a_2), \Phi^+(a_1 + a_2)\} \text{ for } i = 3, 4.$$

Let $i = 3, 4$. Then $\text{Range}(T(a_i, a_1 + a_2)) \subseteq \text{Range}(T(a_i, a_1)) + \text{Range}(T(a_i, a_2))$ so we have $\Phi^+(a_1 + a_2) \in \text{Span}\{\Phi^+(a_i), \Phi^+(a_1), \Phi^+(a_2)\}$. We now use equation (3.2.a) to see

$$(3.2.b) \quad \Phi^+(a_1 + a_2) \in \text{Span}\{\Phi^+(a_1), \Phi^+(a_2)\}.$$

We define

$$\mathcal{A}([a]) := \{a_1 \in A - \{0\} : \Phi([a_1]) \perp \Phi([a])\} \cup \{0\}.$$

It is immediate that $\mathcal{A}([a])$ is closed under scalar multiplication. Let $a_i \in \mathcal{A}([a])$. We wish to show that $a_1 + a_2 \in \mathcal{A}([a])$. This is immediate if $0 = a_1 \wedge a_2$; if $0 \neq a_1 \wedge a_2$, then we use equation (3.2.b) to see $a_1 + a_2 \in \mathcal{A}([a])$. Thus $\mathcal{A}([a])$ is a linear subspace of A . Since Φ is a homeomorphism from $\mathbb{P}(\mathcal{A}([a]))$ to $\mathbb{P}(\Phi^+(a)^\perp)$, we use invariance of domain to see

$$\dim \mathbb{P}(\mathcal{A}([a])) = \dim \mathbb{P}(\Phi^+(a)^\perp) \text{ so } \dim(\mathcal{A}([a])) = q_A - 1.$$

We apply Lemma 2.1 (2) to the family of codimension 1 subspaces $\mathcal{A}([a])$. Set

$$\phi_a(a_1) := T(a, a_1)\Phi^+(a) \text{ for } a_1 \in \mathcal{A}([a]).$$

Since T is bilinear, ϕ_a is linear. If $a_1 \neq 0$, then $\Phi([a_1]) \perp \Phi([a])$ so $0 \neq a \wedge a_1$. By Lemma 2.4 (2), $\{\Phi^+([a]), \phi_a(a_1)\}$ is an orthogonal basis for $\text{Range}(T(a, a_1))$. Since $\Phi([a]) \perp \Phi([a_1])$, $\phi_a(a_1) \in \Phi([a_1])$ so ϕ_a is a linearization of Φ on $\mathcal{A}([a])$. Let $a_i \in A$. Since $q_A \geq 3$ and since Φ is bijective, we may choose $a \in A$ so $\Phi([a]) \perp \Phi([a_1])$ and $\Phi([a]) \perp \Phi([a_2])$. Thus $a_i \in \mathcal{A}([a])$ so every 2 plane is contained in some element of $\mathcal{F}$. Thus by Lemma 2.1 (2) there exists a linearization ϕ of Φ on all of A .

We use equation (1.3.a) to define T_ϕ . Let $0 \neq a_1 \wedge a_2$. Because

$$T_\phi(a_1, a_2) \in \mathfrak{so}(B, \langle \cdot, \cdot \rangle_B) \text{ and } \text{Range}(T_\phi) = \text{Range}(T) = \text{Span}\{\phi(a_1), \phi(a_2)\},$$

we may apply Lemma 2.1 (4) to see $T_\phi(a_1, a_2) = \mu(a_1, a_2)T(a_1, a_2)$. If $0 \neq a_1 \wedge a_2 \wedge a_3$, then $T(a_1, a_2)$ and $T(a_1, a_3)$ are linearly independent maps by Lemma 3.1 (1). We compute:

$$\begin{aligned} T_\phi(a_1, a_2 + a_3) &= \mu(a_1, a_2 + a_3)T(a_1, a_2 + a_3) \\ &= \mu(a_1, a_2 + a_3)T(a_1, a_2) + \mu(a_1, a_2 + a_3)T(a_1, a_3) \\ &= T_\phi(a_1, a_2) + T_\phi(a_1, a_3) \\ &= \mu(a_1, a_2)T(a_1, a_2) + \mu(a_1, a_3)T(a_1, a_3). \end{aligned}$$

This implies that $\mu(a_1, a_2) = \mu(a_1, a_2 + a_3) = \mu(a_1, a_3)$ as $\{T(a_1, a_2), T(a_1, a_3)\}$ is a linearly independent subset of $\mathfrak{so}(B, \langle \cdot, \cdot \rangle_B)$. Consequently $\mu(a_1, a_2)$ is independent of a_2 ; a similar argument shows it is independent of a_1 . We denote this common value by μ and express $T(a_1, a_2) = \mu T_\phi(a_1, a_2)$ if $0 \neq a_1 \wedge a_2$. We use Lemma 2.5 to see $T = \mu T_\phi$ and replace ϕ by $\sqrt{|\mu|}\phi$ to complete the proof. $\square$

§4 EXTENDING LINEARIZATIONS

Let $\mathcal{S}_r$ be the set of all spacelike subspaces of A which have dimension at least r . We have the following inclusions: $\mathcal{S}_{q_A} \subsetneq \dots \subsetneq \mathcal{S}_3$. Let $S \in \mathcal{S}_3$ and $\tilde{S} \in \mathcal{S}_3$. We say that $\{S_i\}_{i \in \mathbb{Z}}$ is a *chain linking S and $\tilde{S}$* if $S = S_i$ for some i , if $\tilde{S} = S_j$ for some j , if $S \cap \tilde{S} \subset S_i \cap S_{i+1}$ for all i , and if $S_i \cap S_{i+1} \in \mathcal{S}_3$ for all i . We omit the proof of the following Lemma in the interests of brevity as the proof is straightforward.

4.1 Lemma. *Let $q_A \geq 5$. Any two elements of $\mathcal{S}_3$ can be linked by a chain.*

If $S \in \mathcal{S}_3$, let T_S be the restriction of T to $S \otimes S$ and let

$$\Phi_S([a]) := \bigcap_{[\tilde{a}] \in \mathbb{P}(S), [\tilde{a}] \neq [a]} \text{Range}(T_S(a, \tilde{a})).$$

4.2 Lemma. *Let $p_B = 0$, let $q_A \geq 5$, and let T be admissible.*

- (1) *If $S_1 \in \mathcal{S}_3$, then $\dim \Phi_{S_1}([a]) = 1$ for all $[a] \in \mathbb{P}(S_1)$.*
- (2) *If $S_1 \in \mathcal{S}_3$, if $S_2 \in \mathcal{S}_3$, and if $S_1 \subseteq S_2$, then $\Phi_{S_2}|_{\mathbb{P}(S_1)} = \Phi_{S_1}$.*
- (3) *There exists $\Phi : \mathbb{P}(A^+) \rightarrow \mathbb{P}(B)$ so $\Phi|_{\mathbb{P}(S)} = \Phi_S$ for all $S \in \mathcal{S}_3$.*
- (4) *Either the map Φ is constant or all the maps Φ_S are injective.*

Proof. Let $S_1 \in \mathcal{S}_3$, let $S_2 \in \mathcal{S}_3$, and let $S_1 \subseteq S_2$. Choose $S_3 \in \mathcal{S}_{q_A}$ so $S_2 \subseteq S_3$. Let $[a] \in \mathbb{P}(S_1)$. We have $\Phi_{S_3}([a]) \subseteq \Phi_{S_2}([a]) \subseteq \Phi_{S_1}([a])$ so by Lemma 3.1 (1,2):

$$1 = \dim\{\Phi_{S_3}([a])\} \leq \dim\{\Phi_{S_2}([a])\} \leq \dim\{\Phi_{S_1}([a])\} = 1.$$

Assertions (1) and (2) follow. Fix $[a] \in \mathbb{P}(A^+)$. Let S and $\tilde{S}$ be any two spaces in $\mathcal{S}_3$ which contain $[a]$. We use Lemma 4.1 to link S and $\tilde{S}$ by a chain $\{S_i\}$. By assertion (2),

$$\Phi_{S_i}([a]) = \Phi_{S_i \cap S_{i+1}}([a]) = \Phi_{S_{i+1}}([a]) \text{ for all } i.$$

Consequently $\Phi_S([a]) = \Phi_{\tilde{S}}([a])$ and $\Phi([a]) := \phi_S([a])$ is well defined and independent of the particular S chosen. This proves assertion (3).

We use Lemma 3.1 (4) to prove assertion (4). Suppose there exists $S \in \mathcal{S}_3$ so Φ_S is not injective. Choose $S_1 \in \mathcal{S}_{q_A}$ so $S \subset S_1$. Then Φ_{S_1} is not injective so Φ_{S_1} and hence Φ_S is constant by Lemma 3.2 (4). Use Lemma 4.2 to link S to any other element $\tilde{S} \in \mathcal{S}_3$ by a chain $\{S_i\}$. If Φ_{S_i} is not injective, then Φ_{S_i} is constant. Thus $\Phi_{S_i \cap S_j}$ is constant and hence $\Phi_{S_{i+1}}$ is constant and takes the same values. Thus Φ_S is not injective implies $\Phi_{\tilde{S}}$ and Φ_S are both constant and take the same value. $\square$

We can now generalize Lemma 3.2 to the case $p_A \neq 0$:

4.3 Lemma. *Let $p_B = 0$ and let $q_A \geq 5$. Let T be admissible.*

- (1) *Let Φ be constant. Fix $\xi \in \mathbb{S}^+(\Phi)$. Set $\chi(a_1, a_2) := -T(a_1, a_2)\xi$. Then $T = T_{\chi, \xi}$.*
- (2) *Let Φ be non-constant. Then there exists a linearization ϕ of Φ so $T = \pm T_\phi$.*

Proof. Suppose that Φ is constant. If a_1 and a_2 span a spacelike 2 plane, then Lemma 3.2 (2) shows that $T(a_1, a_2) = T_{\chi, \xi}(a_1, a_2)$. Lemma 2.5 now implies $T = T_{\chi, \xi}$ which proves assertion (1).

Suppose that Φ is non-constant. We use Lemma 4.2 (4) to see Φ_S is injective for all $S \in \mathcal{S}_3$. As $\mathbb{S}^+(A)$ is the universal cover of $\mathbb{P}(A^+)$, we lift Φ to define

$$\Phi^+ : \mathbb{S}^+(A) \rightarrow \mathbb{S}^+(B) \text{ so } [\Phi^+(a)] = \Phi([a]) \forall a \in \mathbb{S}^+(A).$$

Let $S \in \mathcal{S}_3$. Choose $\bar{S} \in \mathcal{S}_{q_A}$ so $S \subset \bar{S}$. Since $\Phi_{\bar{S}}$ is injective, we apply Lemma 3.2 (2) to T_S to find $\phi_{\bar{S}}$ so $T_{\bar{S}} = T_{\phi_{\bar{S}}}$; we let $\phi_S = \phi_{\bar{S}}|_S$. Then $T_S = T_{\phi_S}$. By Lemma 2.6 (2), ϕ_S is determined up to sign. We use Φ^+ to normalize the choice of sign by requiring

$$(\Phi^+(a), \phi_S(a))_B > 0 \text{ for all } a \in \mathbb{S}^+(S).$$

If $S_1 \in \mathcal{S}_3$, if $S_2 \in \mathcal{S}_3$, and if $S_1 \subset S_2$, we have $\phi_{S_2}|_{S_1} = \phi_{S_1}$. Thus we may use Lemma 4.1 to find

$$\phi : A^+ \rightarrow B \text{ so } \phi|_S = \phi_S \text{ for all } S \in \mathcal{S}_3.$$

We now extend ϕ to all of A . Fix $a \in A$. Let $a_1 \in \mathbb{S}^+(A)$ and $a_2 \in \mathbb{S}^+(A)$ satisfy $a \perp a_i$. Choose $a_3 \in \mathbb{S}^+(A)$ so $\{a, a_1, a_3\}$ and $\{a, a_2, a_3\}$ are orthogonal sets. Choose $\varepsilon_i \in \mathbb{R}$ so that $|\varepsilon_i^2| > |\langle a, a_i \rangle_A|$ for $1 \leq i \leq 3$. As $a + \varepsilon_i a_i \in A^+$, $\phi(a + \varepsilon_i a_i) - \phi(\varepsilon_i a_i)$ is defined. As

$$\begin{aligned} \text{Span}\{a + \varepsilon_1 a_1, a_2, \varepsilon_3 a_3\} &\in \mathcal{S}_3, \text{ Span}\{\varepsilon_1 a_1, a_2, \varepsilon_3 a_3\} \in \mathcal{S}_3, \\ \text{Span}\{a + \varepsilon_3 a_3, \varepsilon_1 a_1, a_2\} &\in \mathcal{S}_3, \end{aligned}$$

and as ϕ_S is linear if $S \in \mathcal{S}_3$, we may compute:

$$\begin{aligned} &\phi(a + \varepsilon_1 a_1) - \phi(\varepsilon_1 a_1) \\ &= \phi(a + \varepsilon_1 a_1 + \varepsilon_3 a_3) - \phi(\varepsilon_3 a_3) - \phi(\varepsilon_1 a_1 + \varepsilon_3 a_3) + \phi(\varepsilon_3 a_3) \\ &= \phi(a + \varepsilon_1 a_1 + \varepsilon_3 a_3) - \phi(\varepsilon_1 a_1) - \phi(\varepsilon_1 a_1 + \varepsilon_3 a_3) + \phi(\varepsilon_1 a_1) \\ &= \phi(a + \varepsilon_3 a_3) - \phi(\varepsilon_3 a_3). \end{aligned}$$

As the roles of a_1 and a_2 are symmetric, this shows that

$$\phi(a + \varepsilon_1 a_1) - \phi(\varepsilon_1 a_1) = \phi(a + \varepsilon_3 a_3) - \phi(\varepsilon_3 a_3) = \phi(a + \varepsilon_2 a_2) - \phi(\varepsilon_2 a_2)$$

is independent of the choice of (a_1, ε_1) . If $a \in A^+$, then this difference yields $\phi(a)$. Consequently we may extend ϕ from A^+ to A by defining

$$\phi(a) := \phi(a + \varepsilon_1 a_1) - \phi(\varepsilon_1 a_1).$$

It is immediate that $\phi(\varrho a) = \varrho \phi(a)$. To check ϕ is linear, we must show ϕ is additive. If $\{a, \tilde{a}\}$ are given, choose $a_1 \in \mathbb{S}^+(A)$ and $\tilde{a}_1 \in \mathbb{S}^+(A)$ so $\{a, a_1, \tilde{a}_1\}$ and $\{\tilde{a}, a_1, \tilde{a}_1\}$ are orthonormal sets. Since $\{a + \tilde{a}, a_1 + \tilde{a}_1\}$ is an orthogonal set and $a_1 + \tilde{a}_1 \in A^+$, we may compute

$$\begin{aligned} \phi(a + \tilde{a}) &= \phi(a + \tilde{a} + \varepsilon(a_1 + \tilde{a}_1)) - \varepsilon \phi(a_1 + \tilde{a}_1) \\ &= \phi(a + \varepsilon a_1) + \phi(\tilde{a} + \varepsilon \tilde{a}_1) - \varepsilon \phi(a_1) - \varepsilon \phi(\tilde{a}_1) \\ &= \phi(a) + \phi(\tilde{a}). \end{aligned}$$

This shows that ϕ is a linear map from A to B . Furthermore, $T(a_1, a_2) = T_\phi(a_1, a_2)$ if $\{a_1, a_2\}$ spans a spacelike 2 plane. We use Lemma 2.5 to see $T = T_\phi$. $\square$

§5 THE PROOF OF THEOREMS 1.2 (2) AND 1.4 (2)

5.1 The proof of Theorem 1.4 (2) for metrics of arbitrary signature. Let $q_A \geq 5$. We use Lemma 2.2 (2) to find ψ which relates $\langle \cdot, \cdot \rangle_B$ to a positive definite metric $\langle \cdot, \cdot \rangle_+$. Let T be admissible. Then ψT is admissible and takes values in $\mathfrak{so}(B, \langle \cdot, \cdot \rangle_+)$. We use Lemma 4.3 to see

$$(5.1.a) \quad \begin{aligned} \psi T(a_1, a_2)b &= (\hat{\chi}(a_1, a_2), b)_+ \hat{\xi} - (\hat{\xi}, b)_+ \hat{\chi}(a_1, a_2) \text{ or} \\ \psi T(a_1, a_2)b &= \pm \{(\hat{\phi}(a_2), b)_+ \hat{\phi}(a_1) - (\hat{\phi}(a_1), b)_+ \hat{\phi}(a_2)\}. \end{aligned}$$

Since $\psi^2 = id$, we may apply ψ to equation (5.1.a) and use Lemma 2.2 (1) to see:

$$\begin{aligned} T(a_1, a_2)b &= \langle \psi \hat{\chi}(a_1, a_2), b \rangle_B \psi \hat{\xi} - \langle \psi \hat{\xi}, b \rangle_B \psi \hat{\chi}(a_1, a_2) \text{ or} \\ T(a_1, a_2)b &= \pm \{ \langle \psi \hat{\phi}(a_2), b \rangle_B \psi \hat{\phi}(a_1) - \langle \psi \hat{\phi}(a_1), b \rangle_B \psi \hat{\phi}(a_2) \}. \end{aligned}$$

We complete the proof by setting $(\chi, \xi) := (\psi \hat{\chi}, \psi \hat{\xi})$ or $\phi := \psi \hat{\phi}$ as appropriate. $\square$

5.2 The proof of Theorem 1.2 (2). Let $\langle \cdot, \cdot \rangle$ be a non-degenerate metric of signature (p, q) on V where $q \geq 5$. Let R be an algebraic curvature tensor which has constant rank 2. We apply Theorem 1.4 (2). Suppose first that $R = R_{\chi, \xi}$. Choose $\{x, y\}$ a spacelike orthonormal set so $\{\xi, x, y\}$ is an orthogonal set. Choose z so $\langle \xi, z \rangle \neq 0$. We apply the Bianchi identity (1.2) to see:

$$\begin{aligned} 0 &= \{ \langle \chi(x, y), z \rangle + \langle \chi(y, z), x \rangle + \langle \chi(z, x), y \rangle \} \xi \\ &\quad - \langle \xi, z \rangle \chi(x, y) - \langle \xi, x \rangle \chi(y, z) - \langle \xi, y \rangle \chi(z, x). \end{aligned}$$

Since $\langle \xi, z \rangle \neq 0$ but $\langle \xi, x \rangle = \langle \xi, y \rangle = 0$ we see $\chi(x, y)$ is a multiple of ξ so

$$\dim\{\text{Range}(R(x, y))\} \leq 1$$

which is false. Thus $R = \pm R_\phi$ where $\ker(\phi) \cap V^+ = \emptyset$. By replacing R by $-R$ we may suppose $R = R_\phi$. Since R_ϕ is an algebraic curvature tensor, we may apply Lemma 2.3 (2) to see $\phi = \phi^*$. $\square$

§6 THE CLASSIFICATION OF IVANOV-PETROVA ALGEBRAIC CURVATURE TENSORS OF RANK 2

Let R be a rank 2 algebraic curvature tensor of rank 2. We use Theorem 1.2 to express $R = \pm R_\phi$ where ϕ is admissible. Let $a_1 \in \mathbb{S}^+(V)$. Choose $S \in \mathcal{S}_3(V)$ so $a_1 \in S$. Choose $a_2 \in \mathbb{S}^+(S)$ so $a_2 \perp a_1$ and so $\phi(a_2) \perp \phi(a_1)$. Let $\pi := \text{Span}\{a_1, a_2\}$ and let $\sigma := \text{Range}(R_\phi(\pi))$. Since $\{\phi(a_1), \phi(a_2)\}$ is an orthogonal basis for σ , we have

$$R_\phi(\pi) : \phi(a_1) \rightarrow -\langle \phi(a_1), \phi(a_1) \rangle \phi(a_2) \text{ and } R_\phi(\pi) : \phi(a_2) \rightarrow \langle \phi(a_2), \phi(a_2) \rangle \phi(a_1).$$

Let $\langle \cdot, \cdot \rangle_\sigma$ be the restriction of $\langle \cdot, \cdot \rangle$ to σ . Since $\langle \phi(a_1), \phi(a_2) \rangle = 0$, $\langle \cdot, \cdot \rangle_\sigma$ is determined by the two inner products $\langle \phi(a_1), \phi(a_1) \rangle$ and $\langle \phi(a_2), \phi(a_2) \rangle$. We can relate the eigenvalue structure of $R_\phi(\pi)$ to $\langle \cdot, \cdot \rangle_\sigma$, there are 5 cases:

- (1) The inner product $\langle \cdot, \cdot \rangle_\sigma$ is positive or negative definite. Then $R_\phi(\pi)$ has two non-zero purely imaginary eigenvalues $\pm \sqrt{-1} \lambda$ where $\lambda^2 = \langle \phi(a_1), \phi(a_1) \rangle \langle \phi(a_2), \phi(a_2) \rangle$.
- (2) The inner product $\langle \cdot, \cdot \rangle_\sigma$ is indefinite and non-degenerate. Then $R_\phi(\pi)$ has two non-zero real eigenvalues $\pm \lambda$ where $\lambda^2 = -\langle \phi(a_1), \phi(a_1) \rangle \langle \phi(a_2), \phi(a_2) \rangle$.
- (3) The inner product $\langle \cdot, \cdot \rangle_\sigma$ is degenerate but non-trivial. Then $R_\phi^2 \neq 0$ but $R_\phi^3 = 0$.
- (4) The inner product $\langle \cdot, \cdot \rangle_\sigma$ is trivial. Then $R_\phi^2 = 0$.

We can now prove one implication of Theorem 1.3. Suppose either that there exists $C \neq 0$ so that $\langle \phi(v_1), \phi(v_2) \rangle = C \langle v_1, v_2 \rangle$ for all v_i or that $\langle \phi(v_1), \phi(v_2) \rangle = 0$ for all v_i . Then either case (1) or case (4) holds and R is Ivanov-Petrova.

To prove the other implication of Theorem 1.3, we suppose that R_ϕ is Ivanov-Petrova. We first show that the complex Jordan normal form described by cases (2) and (3) can not hold. Let $v_1 \in \mathbb{S}(V^+)$. Since $q \geq 5$, we may choose $S \in \mathcal{S}_5$ so that $v_1 \in S$. Choose $v_2 \in S$ as above. Choose $v_3 \in S$ so $v_3 \perp v_i$ and $\phi(v_3) \perp \phi(v_i)$ for $i = 1, 2$. Then $\{v_1, v_2, v_3\}$ is a spacelike orthonormal set and $\{\phi(v_1), \phi(v_2), \phi(v_3)\}$ is an orthogonal linearly independent set. For $1 \leq i < j \leq 3$, let $\pi_{ij} := \text{Span}\{v_i, v_j\}$. Suppose $R_\phi(\pi_{ij})$ has two non-zero real eigenvalues. Then:

$$\langle \phi(v_i), \phi(v_i) \rangle \cdot \langle \phi(v_j), \phi(v_j) \rangle < 0 \text{ for } 1 \leq i < j \leq 3.$$

This is not possible as at least two of the inner products must have the same sign. Suppose $R_\phi(\pi_{ij})^3 = 0$ but $R_\phi(\pi_{ij})^2 \neq 0$. Then the set of inner products

$$\{\langle \phi(v_i), \phi(v_i) \rangle, \langle \phi(v_j), \phi(v_j) \rangle\}$$

consists of one zero number and one non-zero number for $1 \leq i < j \leq 3$. Again, this is not possible as there are 3 such pairs.

We now deal with the remaining eigenvalue structures. Suppose $R_\phi(\pi)$ has two non-zero purely imaginary eigenvalues $\pm \sqrt{-1}\lambda$ for all oriented spacelike 2 planes π . Let v_1 and v_2 be arbitrary unit spacelike vectors in V . Since $q \geq 5$, we can choose a third unit spacelike vector v_3 so $v_3 \perp v_1$, $v_3 \perp v_2$, $\phi(v_3) \perp \phi(v_1)$, and so $\phi(v_3) \perp \phi(v_2)$. Let $\pi_i := \text{Span}\{v_i, v_3\}$ for $i = 1, 2$. We then have

$$0 \neq \lambda^2 = \langle \phi(v_1), \phi(v_1) \rangle \cdot \langle \phi(v_3), \phi(v_3) \rangle = \langle \phi(v_2), \phi(v_2) \rangle \cdot \langle \phi(v_3), \phi(v_3) \rangle \text{ so} \\ \langle \phi(v_1), \phi(v_1) \rangle = \langle \phi(v_2), \phi(v_2) \rangle = C \text{ for all } v_i \in \mathbb{S}(V^+) \text{ where } C = \pm \lambda.$$

We rescale this identity to see that $\langle \phi(v), \phi(v) \rangle = C \langle v, v \rangle$ for all spacelike v . Now let v be arbitrary. Choose v_1 spacelike. Then $v + \varepsilon v_1$ is spacelike for large ε and hence

$$\langle \phi(v + \varepsilon v_1), \phi(v + \varepsilon v_1) \rangle = C \langle v + \varepsilon v_1, v + \varepsilon v_1 \rangle \text{ for } \varepsilon > \varepsilon_0.$$

Since this equation is quadratic in the parameter ε , it must hold $\varepsilon = 0$ so

$$\langle \phi(v), \phi(v) \rangle = C \langle v, v \rangle \text{ for all } v \in V.$$

Suppose that $R_\phi(\pi)^2 = 0$ for any spacelike 2 plane π . Then $\text{Range}(R_\phi(\pi))$ is totally isotropic. Let v_1 and v_2 be arbitrary (not necessarily distinct) vectors in V . Choose unit spacelike vectors w_1 and w_2 so $w_1 \perp w_2$. Let $v_i(\varepsilon) := v_i + \varepsilon w_i$. For large ε , $\text{Span}\{v_1(\varepsilon), v_2(\varepsilon)\}$ is spacelike so $\langle \phi(v_1(\varepsilon)), \phi(v_2(\varepsilon)) \rangle = 0$. Since this identity is quadratic, it continues to hold for $\varepsilon = 0$ and thus $\langle \phi(v_1), \phi(v_2) \rangle = 0$ for all $v_i \in V$ so $\text{Range}(\phi)$ is totally isotropic. This completes the proof of Theorem 1.3.

§7 REALIZING RANK 2 ALGEBRAIC CURVATURE TENSORS

Let W be a vector space with a metric of signature (r, s) and let $F : M \rightarrow W$ be an immersion of a manifold M into W as a non-degenerate hypersurface; this means that the first fundamental form is a non-degenerate metric of signature (p, q) on M where $(p, q) = (r - 1, s)$ or $(p, q) = (r, s - 1)$. Choose a normal ν along M so $\langle \nu, \nu \rangle_W = \pm 1$ and so $\langle dF, \nu \rangle_W = 0$. Let L be the second fundamental form and let S be the associated shape operator determined by the immersion. These two tensors are related by the identity:

$$L(v_1, v_2) = \langle v_1 v_2(F), \nu \rangle_W = \langle S v_1, v_2 \rangle_W.$$

7.1 Lemma. *If $(p, q) = (r - 1, s)$, then $R = -R_S$; if $(p, q) = (r, s - 1)$, then $R = R_S$.*

Proof. Although this is well known, we sketch the proof to establish later notation. Fix a point $P \in M$. Choose a basis $\{e_1, \dots, e_{m+1}\}$ for W so that $\{e_1, \dots, e_m\}$ is a basis for $T_P M$ and so that $e_{m+1} = \nu(P)$. Let $\{x^1, \dots, x^{m+1}\}$ be the dual basis for W^* ; these give coordinates on W . Then $\vec{y} := (F^* x^1, \dots, F^* x^m)$ is a system of local coordinates on M and the immersion takes the form $F(\vec{y}) = y^1 e_1 + \dots + y^m e_m + f(\vec{y}) e_{m+1}$ where f is a smooth scalar valued function with $df(P) = 0$. Let $\varepsilon = \langle e_{m+1}, e_{m+1} \rangle_W = \pm 1$. We complete the proof by computing:

$$\begin{aligned} \langle S \partial_i, \partial_j \rangle(P) &= L(\partial_i, \partial_j)(P) = \langle \partial_i \partial_j F, e_{m+1} \rangle_W(P) = \varepsilon_{m+1} \partial_i \partial_j f(P), \\ g_{ij} &= \langle e_i, e_j \rangle_W + \varepsilon \partial_i(f) \partial_j(f), \quad \partial_k(g_{ij})(P) = 0, \\ R_{ijkl}(P) &= \frac{1}{2} \{ \partial_i \partial_k g_{jl} + \partial_j \partial_l g_{ik} - \partial_i \partial_l g_{jk} - \partial_j \partial_k g_{il} \}(P) \\ &= \varepsilon \{ \partial_i \partial_l(f) \cdot \partial_j \partial_k(f) - \partial_i \partial_k(f) \cdot \partial_j \partial_l(f) \}(P), \\ (R_S)_{ijkl}(P) &= \{ \langle S \partial_i, \partial_l \rangle \langle S \partial_j, \partial_k \rangle - \langle S \partial_i, \partial_k \rangle \langle S \partial_j, \partial_l \rangle \}(P) = \varepsilon R_{ijkl}(P). \quad \square \end{aligned}$$

Proof of Theorem 1.5. By Theorem 1.2, we may choose ϕ so $R = \varepsilon R_\phi$ where $\varepsilon = \pm 1$; ϕ is symmetric and $\ker(\phi)$ contains no spacelike vectors. We adopt the notation of equation (1.4.a) to define the germ of an immersion F_ϕ with associated first fundamental form g_ϕ , second fundamental form L_ϕ , and shape operator S_ϕ . If $\{e_i\}$ is a basis for V with associated coordinates $\{y^i\}$, then

$$\begin{aligned} F_\phi(\vec{y}) &:= y^1 e_1 + \dots + y^m e_m + \frac{1}{2} \sum_{i,j} y^i y^j \langle \phi(e_i), e_j \rangle_V \cdot e_{m+1}, \\ \partial_i(F_\phi) &= e_i + \sum_k y^k \langle \phi(e_i), e_k \rangle_V \cdot e_{m+1}, \text{ and} \\ g_{ij}(\vec{y}) &= \langle e_i, e_j \rangle_V + \varepsilon \sum_{k,l} y^k y^l \langle \phi(e_i), e_k \rangle_V \cdot \langle \phi(e_j), e_l \rangle_V. \end{aligned}$$

This shows that $g_{ij}(0) = \langle \cdot, \cdot \rangle_V$. Since e_{m+1} is the normal at $\vec{y} = 0$, we have that $L_\phi(0)(e_i, e_j) = \varepsilon \langle \phi(e_i), e_j \rangle_V$. Thus $S_\phi(0) = \varepsilon \phi$ so by Lemma 7.1, ${}^g R = \varepsilon R_{\varepsilon \phi} = R$. Consequently (V, g_ϕ) is a geometric realization of R .

We may choose the basis $\{e_i\}$ so that $\{e_i\}_{i>s}$ is a basis for $\ker(\phi)$; if ϕ is invertible, then $s = m$ and this is the empty basis. If $i > s$, we have $\partial_i \partial_j(F) = \langle \phi(e_i), e_j \rangle_V = 0$ so $L_\phi(P)(\partial_i, \partial_j) = 0$ and $S_\phi(P)(\partial_i) = 0$ at any point P . Since we are only considering an arbitrarily small neighborhood of the origin, $\text{Rank}(S_\phi(P)) \geq \text{Rank}(S_\phi(0)) = s$. Thus equality holds and $\ker(S_\phi(P)) = \text{Span}\{e_i\}_{i>s}$. If $i, j > s$, then $g_{ij}(P) = g_{ij}(0)$. Since $\ker(\phi) = \text{Span}\{e_i\}_{i>s}$ contains no spacelike vectors, $\ker(S_\phi)(P)$ contains no spacelike vectors and (V, g_ϕ) is the germ of a rank 2 pseudo-Riemannian manifold. $\square$

§8 EXAMPLES

8.1 Example. We have assumed in the previous sections that $q_A \geq 5$. We now show that Theorem 1.4 (2) fails if $q_A = 4$. Let $\{e_1^-, \dots, e_p^-, e_1^+, \dots, e_q^+\}$ be a basis for $\mathbb{R}^{(p,q)}$. We define $\langle \cdot, \cdot \rangle_{(p,q)}$ by:

$$\langle e_i^-, e_j^- \rangle_{(p,q)} = -\delta_{ij}, \quad \langle e_i^-, e_j^+ \rangle_{(p,q)} = 0, \quad \text{and} \quad \langle e_i^+, e_j^+ \rangle_{(p,q)} = \delta_{ij}.$$

We suppose first $p_A = 0$. Let $\text{orn} := e_1^+ \wedge e_2^+ \wedge e_3^+ \wedge e_4^+$ be the standard orientation of $\mathbb{R}^{(0,4)}$. Extend the inner product on $\mathbb{R}^{(0,4)}$ to an inner product on $\Lambda^4(\mathbb{R}^{(0,4)})$ and define:

$$T^{(0,4)} : \mathbb{R}^{(0,4)} \otimes \mathbb{R}^{(0,4)} \rightarrow \mathfrak{so}(\mathbb{R}^{(0,4)}, \langle \cdot, \cdot \rangle_{(0,4)}) \text{ by}$$

$$\langle T^{(0,4)}(v_1, v_2)v_3, v_4 \rangle_{(0,4)} = \langle v_1 \wedge v_2 \wedge v_3 \wedge v_4, \text{orn} \rangle_{(0,4)}.$$

Let $\{f_1, f_2\}$ be an oriented orthonormal basis for a 2 plane π . We extend $\{f_1, f_2\}$ to an oriented orthonormal basis $\{f_1, f_2, f_3, f_4\}$ for $\mathbb{R}^{(0,4)}$ and compute:

$$T^{(0,4)}(f_1, f_2)f_1 = 0, \quad T^{(0,4)}(f_1, f_2)f_2 = 0,$$

$$T^{(0,4)}(f_1, f_2)f_3 = f_4, \quad \text{and} \quad T^{(0,4)}(f_1, f_2)f_4 = -f_3.$$

Thus $T^{(0,4)}$ has constant rank 2. Since $\text{Range}(T^{(0,4)}(\pi)) = \pi^\perp$, $\Phi([a]) = \{0\}$. Thus $T^{(0,4)} \neq T_\phi^{(0,4)}$ and $T^{(0,4)} \neq T_{\chi, \xi}^{(0,4)}$ for any ϕ or (χ, ξ) . More generally, let $\sigma : \mathbb{R}^{(p,4)} \rightarrow \mathbb{R}^{(0,4)}$ denote projection on the last 4 components. We extend $T^{(0,4)}$ to $\mathbb{R}^{(p,4)} \otimes \mathbb{R}^{(p,4)}$ and thereby construct an algebraic curvature tensor which does not satisfy the conclusions of Theorem 1.4 (2) for any p by defining:

$$T^{(p,4)}(v_1, v_2) := T^{(0,4)}(\sigma(v_1), \sigma(v_2)).$$

8.2 Example. Let $q = p + 1 \geq 5$. We define

$$\phi(e_i^\pm) = \pm(e_i^- + e_i^+) \text{ for } i \leq p \text{ and } \phi(e_{p+1}^+) = e_{p+1}^+.$$

Then ϕ is self-adjoint and $\ker(\phi) = \text{Span}\{e_i^- - e_i^+\}_{i \leq p}$ contains no spacelike vectors so R_ϕ has constant rank 2. Let π be a spacelike 2 plane. Since the metric on $\text{Range}(R(\pi))$ is degenerate, $R_\phi^3(\pi) = 0$ so 0 is the only eigenvalue of R_ϕ . Let $\pi_i := \text{Span}\{e_1^+, e_i^+\}$. Then $R_\phi(\pi_2)^2 = 0$ and $R_\phi(\pi_{p+1})^2 \neq 0$ so R_ϕ is not an Ivanov-Petrova algebraic curvature tensor. When dealing with an indefinite metric, it is necessary to study the complex Jordan normal form and not just the eigenvalue structure.

8.3 Example. We now construct an algebraic curvature tensor which is (spacelike) rank 2 Ivanov-Petrova but which does not have constant rank on the Grassmanian of oriented timelike 2 planes as follows. Let $q = p - 1 \geq 5$. We define:

$$\phi(e_i^\pm) = \pm(e_i^- + e_i^+) \text{ for } i \leq q, \text{ and } \phi(e_{q+1}^-) = 0.$$

Since ϕ is self-adjoint and $\ker(\phi) = \text{Span}\{e_i^- + e_i^+, e_{q+1}^-\}_{i \leq q}$ contains no spacelike vectors, $R_\phi(\pi)$ has rank 2. Furthermore as the range of ϕ is totally isotropic, R_ϕ is (spacelike) rank 2 Ivanov-Petrova. Let $\pi_i^- := \text{Span}\{e_1^-, e_i^-\}$ be oriented timelike 2 planes. Then $\text{Rank}(R_\phi(\pi_i)) = 2$ if $2 \leq i \leq q$ while $\text{Rank}(R_\phi(\pi_i)) = 0$ if $i = q + 1$. Thus the rank of R_ϕ is not constant on the timelike 2 planes.

Acknowledgements. The research of Gilkey was partially supported by the NSF (USA) and MPI (Leipzig, Germany). It is a pleasant task to thank Professors Cortes, Simon, and Vrancken for helpful discussions concerning the material of §7.

REFERENCES

- N. Blažić, N. Bokan and P. Gilkey, *A Note on Osserman Lorentzian manifolds*, Bull. London Math. Soc. **29** (1997), 227–230.
- N. Blažić, N. Bokan, P. Gilkey, and Z. Rakić, *Pseudo-Riemannian Osserman manifolds*, J. Balkan Society of Geometers **2** (1997), 1–12.
- Q.-S. Chi, *A curvature characterization of certain locally rank-one symmetric spaces*, J. Differential Geom. **28** (1988), 187–202.
- E. García-Riό, D. Kupeli, and M. Vázquez-Abal, *On a problem of Osserman in Lorentzian geometry*, Differential Geom. Appl. **7** (1997), 85–100.
- E. García-Riό, M. E. Vázquez-Abal and R. Vázquez-Lorenzo, *Nonsymmetric Osserman pseudo-Riemannian manifolds*, Proc. Amer. Math. Soc. **126** (1998), 2763–2769.
- P. Gilkey, *Geometric properties of the curvature operator*, To appear, Geometry and Topology of Submanifolds **X**, World Scientific.
- , *Manifolds whose curvature operator has constant eigenvalues at the basepoint*, J. Geom. Anal. **4** (1994), 155–158.
- , *Riemannian manifolds whose skew-symmetric curvature operator has constant eigenvalues II*, in **Differential geometry and applications** (eds Kolar, Kowalski, Krupka, and Slovák) Publ Massaryk University Brno Czech Republic ISBN 80-210-2097-0 (1999), 73–87.
- , *Natural Operators Defined by the Riemann Curvature Tensor*, to appear Fall 2001, World Scientific.
- P. Gilkey, J. V. Leahy and H. Sadofsky, *Riemannian manifolds whose skew-symmetric curvature operator has constant eigenvalues*, Indiana Univ. Math. J. **48** (1999), 615–634.
- P. Gilkey and U. Semmelman, *Spinors, self-duality, and IP algebraic curvature tensors of rank 4*, preprint (1999).
- S. Ivanov and I. Petrova, *Riemannian manifold in which the skew-symmetric curvature operator has pointwise constant eigenvalues*, Geom. Dedicata **70** (1998), 269–282.
- O. Kowalski, M. Sekizawa, and Z. Vlášek, *Can tangent sphere bundles over Riemannian manifolds have strictly positive sectional curvature?*, preprint.
- R. Osserman, *Curvature in the eighties*, Amer. Math. Monthly **97** (1990), 731–756.
- T. Zhang, *Manifolds with indefinite metrics whose skew-symmetric curvature operator has constant eigenvalues*, Ph. D. thesis, University of Oregon (2000).

PG: Mathematics Department, University of Oregon, Eugene Or 97403 USA
E-mail address: gilkey@darkwing.uoregon.edu

TZ: Mathematics Department, Murray State University, Murray Kentucky 42071 USA
E-mail address: tan.zhang@murraystate.edu